

Estrogen/Progesterone - Thaw Cycle

Introduction

Estrogen and progesterone can be used to supplement the natural cycle to time embryo replacement of frozen/thawed embryos

The protocol described herein is called the "EEP" protocol for <u>E</u>xogenous <u>E</u>strogen and <u>P</u>rogesterone

Before Cycle Start

- Start taking a Prenatal Vitamin.
- Review questions with doctor
- Consent forms signed and witnessed at Boston IVF
- ♦ Approval from financial dept.

Cycle Start

- Call IVF Nurse on cycle day one -
- On DAY ONE, start Estrace tablets, each 2 mg, and take one tablet twice daily
- On DAY ONE, start Estradiol patch, apply first patch and

change every 3rd day or as instructed by your nurse.

Start progesterone at the instruction of your nurse. Embryo transfer day 4 of progesterone, day 6 if Blastocyst.

- Continue with Estrace/Estradiol patch and Progesterone until the pregnancy test is done.
- Blood tests and ultrasound examinations are determined by your physician

Embryo Transfer

- Preformed on day 4 or day 6 of progesterone.
- The IVF Nurse will call you two days prior to embryo transfer to determine what time to come in.
- Simple procedure, no anesthesia, home after 10 minutes in recovery room.

 On the day of the transfer take it easy, no heavy lifting over 20lbs and no formal exercise. After the transfer you may resume regular activities.

Pregnancy Test

- Performed 12-10 days after the embryo transfer
- Pregnancy test will be scheduled by the nurse in the recovery room.

Good luck and best wishes from all of us at Boston IVF!

